

Neutrino masses and mixing from neutrino oscillation data

S.M. Bilenky^a, C. Giunti^b and W. Grimus^c

^aJoint Institute for Nuclear Research, Dubna, Russia

^bINFN, Sezione di Torino, and Dipartimento di Fisica Teorica, Via P. Giuria 1, I-10125 Torino, Italy

^cInstitute for Theoretical Physics, University of Vienna, Boltzmanngasse 5, A-1090 Vienna, Austria

We discuss what information about the neutrino mass spectrum and the elements of the neutrino mixing matrix can be inferred from the results of neutrino oscillation experiments. The cases of three and four massive neutrinos are considered. It is shown that the general characteristics of the neutrino mixing matrix are quite different from those of the quark mixing matrix. Talk presented by S.M. Bilenky at the XVI *International Workshop on Weak Interactions and Neutrinos*, Capri, Italy, June 22–28 1997. Preprint DFTT 45/97, hep-ph/9707358.

The problem of neutrino masses and mixing is the central issue of today's neutrino physics. The results of many experiments on the search for effects of neutrino masses and mixing were discussed at this meeting. New and more precise experiments are going on or are under preparation.

We will present here some results of Refs.[1–4], in which model independent information about the spectrum of neutrino masses and the elements of the neutrino mixing matrix was obtained from the existing neutrino oscillation data. We will consider here also possible implications of our results for the future experiments.

In accordance with the neutrino mixing hypothesis [5], the left-handed flavor neutrino fields $\nu_{\alpha L}$ determined by the standard CC and NC interactions are mixtures of the left-handed components of the fields ν_i of neutrinos (Dirac or Majorana) with mass m_i :

$$\nu_{\alpha L} = \sum_{i=1}^n U_{\alpha i} \nu_{iL}, \quad (1)$$

where $U^\dagger U = 1$, $\alpha = e, \mu, \tau, \dots$ and $n \geq 3$.

From the existing neutrino oscillation data it follows that three different scales of neutrino mass-squared differences can be relevant for neutrino oscillations: $\simeq 10^{-5} \text{ eV}^2$ for solar neutrinos [6], $\simeq 10^{-2} \text{ eV}^2$ for atmospheric neutrinos [7] and $\simeq 1 \text{ eV}^2$ for LSND neutrinos [8]. Let us con-

sider the general case of n neutrinos ($m_1 < m_2 < \dots < m_n$) with the largest mass-squared difference $\Delta m^2 \equiv m_n^2 - m_1^2$ relevant for short-baseline (SBL) oscillations and two groups of close masses $m_1 < \dots < m_{r-1}$ and $m_r < \dots < m_n$ such that

$$\begin{aligned} \Delta m_{i1}^2 &\ll \Delta m^2 \quad \text{for } i < r, \\ \Delta m_{nj}^2 &\ll \Delta m^2 \quad \text{for } j \geq r, \end{aligned} \quad (2)$$

with $\Delta m_{ij}^2 \equiv m_i^2 - m_j^2$. In this case, for the SBL transition probabilities we have [2]

$$P_{\nu_\alpha \rightarrow \nu_\beta}^{(\text{SBL})} = \frac{A_{\alpha;\beta}}{2} \left(1 - \cos \frac{\Delta m^2 L}{2p} \right) \quad (\beta \neq \alpha), \quad (3)$$

$$P_{\nu_\alpha \rightarrow \nu_\alpha}^{(\text{SBL})} = 1 - \frac{B_{\alpha;\alpha}}{2} \left(1 - \cos \frac{\Delta m^2 L}{2p} \right), \quad (4)$$

where p is the neutrino momentum and L is the distance between the neutrino source and detector. The oscillation amplitudes are given by

$$A_{\alpha;\beta} = 4 \left| \sum_{i=r}^n U_{\beta i} U_{\alpha i}^* \right|^2, \quad (5)$$

$$B_{\alpha;\alpha} = 4 \left(\sum_{i=r}^n |U_{\alpha i}|^2 \right) \left(1 - \sum_{i=r}^n |U_{\alpha i}|^2 \right). \quad (6)$$

No indication in favour of neutrino oscillations was found in SBL reactor and accelerator disappearance experiments. From the exclusion plots obtained from the results of these experiments it follows that

$$B_{e;e} \leq B_{e;e}^0 \quad \text{and} \quad B_{\mu;\mu} \leq B_{\mu;\mu}^0. \quad (7)$$

The numerical values of the upper bounds $B_{\alpha;\alpha}^0$ ($\alpha = e, \mu$) depend on the value of Δm^2 . We considered the wide interval

$$10^{-1} \leq \Delta m^2 \leq 10^3 \text{ eV}^2. \quad (8)$$

From Eqs.(6) and (7) it follows that

$$\sum_{i=r}^n |U_{\alpha i}|^2 \leq a_{\alpha}^0 \quad \text{or} \quad \sum_{i=r}^n |U_{\alpha i}|^2 \geq 1 - a_{\alpha}^0, \quad (9)$$

with $\alpha = e, \mu$ and

$$a_{\alpha}^0 = \frac{1}{2} \left(1 - \sqrt{1 - B_{\alpha;\alpha}^0} \right). \quad (10)$$

From the results of the Bugey $\bar{\nu}_e \rightarrow \bar{\nu}_e$ [9] and the CDHS and CCFR [10] $\bar{\nu}_{\mu} \rightarrow \bar{\nu}_{\mu}$ experiments it follows that the values of a_e^0 and a_{μ}^0 are small ($a_e^0 \lesssim 4 \times 10^{-2}$ for any value of Δm^2 in the interval (8) and $a_{\mu}^0 \lesssim 2 \times 10^{-1}$ for $\Delta m^2 \gtrsim 0.3 \text{ eV}^2$). Thus, from the existing inclusive data it follows that the sums $\sum_{i=r}^n |U_{ei}|^2$ and $\sum_{i=r}^n |U_{\mu i}|^2$ can be either small or large (i.e., close to one).

We will consider first the simplest case of three massive neutrinos with the mass hierarchy $m_1 \ll m_2 \ll m_3$ and will assume that Δm_{21}^2 is relevant for the suppression of solar ν_e 's. In this case, we have

$$|U_{\alpha 3}|^2 \leq a_{\alpha}^0 \quad \text{or} \quad |U_{\alpha 3}|^2 \geq 1 - a_{\alpha}^0, \quad (11)$$

with $\alpha = e, \mu$. In the case under consideration, the probability of solar ν_e 's to survive is given by [11]

$$P_{\nu_e \rightarrow \nu_e}^{\text{sun}} = |U_{e3}|^4 + (1 - |U_{e3}|^2)^2 P_{\nu_e \rightarrow \nu_e}^{(1;2)}, \quad (12)$$

where $P_{\nu_e \rightarrow \nu_e}^{(1;2)}$ is the probability of ν_e 's to survive due to the coupling of ν_e with ν_1 and ν_2 . From Eq.(12) it follows that, in order to explain the solar neutrino data [6], from the two possibilities (11) for $|U_{e3}|^2$ we must choose $|U_{e3}|^2 \leq a_e^0$. Hence, we come to the two possible schemes:

$$\text{(I)} \quad |U_{e3}|^2 \leq a_e^0, \quad |U_{\mu 3}|^2 \leq a_{\mu}^0; \quad (13)$$

$$\text{(II)} \quad |U_{e3}|^2 \leq a_e^0, \quad |U_{\mu 3}|^2 \geq 1 - a_{\mu}^0. \quad (14)$$

The most natural scheme I, with the hierarchy of couplings corresponding to the hierarchy of neutrino masses, is not favoured by the results of the




Figure 1

LSND experiment. In fact, from Eqs.(5) and (13) we have the following upper bound for the amplitude $A_{\mu;e} = 4|U_{e3}|^2|U_{\mu 3}|^2$ of $\bar{\nu}_{\mu} \rightarrow \bar{\nu}_e$ transitions:

$$A_{\mu;e} \leq 4a_e^0 a_{\mu}^0. \quad (15)$$

As it is seen from Fig.1, the LSND-allowed region that is not excluded by the data of other experiments lies inside of the region forbidden by the inequality (15) (the curve passing through the circles).

On the other hand, scheme II allows to describe the results of all experiments, including LSND. Note that if this scheme is realized in nature we have $m_{\nu_{\mu}} \gg m_{\nu_e}, m_{\nu_{\tau}}$.

Up to now we did not take into account the atmospheric neutrino anomaly [7]. In order to take into account all data we will consider schemes with four massive neutrinos [2–4]. Let us first assume that there is a hierarchy of neutrino masses,

$m_1 \ll m_2 \ll m_3 \ll m_4$, with Δm_{21}^2 relevant for the suppression of solar ν_e 's, Δm_{31}^2 relevant for the atmospheric neutrino anomaly and $\Delta m^2 \equiv \Delta m_{41}^2$ relevant for the LSND anomaly. From the data of inclusive experiments it follows that

$$|U_{\alpha 4}|^2 \leq a_\alpha^0 \quad \text{or} \quad |U_{\alpha 4}|^2 \geq 1 - a_\alpha^0, \quad (16)$$

with $\alpha = e, \mu$. The survival probability of solar ν_e 's is given by Eq.(12) with the obvious change $|U_{e3}| \rightarrow |U_{e4}|$. For the survival probability of atmospheric $\bar{\nu}_\mu$'s we have the lower bound

$$R_{\bar{\nu}_\mu \rightarrow \bar{\nu}_\mu}^{(-)} \geq |U_{\mu 4}|^2. \quad (17)$$

Thus, from the solar and atmospheric neutrino data it follows that both elements $|U_{e4}|^2$ and $|U_{\mu 4}|^2$ are small:

$$|U_{e4}|^2 \leq a_e^0 \quad \text{and} \quad |U_{\mu 4}|^2 \leq a_\mu^0. \quad (18)$$

Hence, for the amplitude $A_{\mu;e} = 4|U_{e4}|^2|U_{\mu 4}|^2$ of $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ transitions we have the upper bound (15) that is presented in Fig.1 (the curve passing through the circles). From the figure one can see that this limit is in contradiction with the LSND-allowed region. Thus, we come to the conclusion that the hierarchy of masses of four neutrinos is not favoured by the results of the LSND and other neutrino oscillation experiments. The same conclusion is valid for all the possible neutrino mass spectra with one mass separated from the group of three close masses by a gap of about 1 eV.

Only the following two mass spectrum of four neutrinos are favoured by all the existing neutrino oscillation data:

$$(A) \quad \underbrace{m_1 < m_2}_{\text{atm}} \ll \underbrace{m_3 < m_4}_{\text{solar}}, \quad (19)$$

LSND

$$(B) \quad \underbrace{m_1 < m_2}_{\text{solar}} \ll \underbrace{m_3 < m_4}_{\text{atm}}. \quad (20)$$

LSND

Taking into account the data of the solar and atmospheric neutrino experiments, for the schemes A and B we have respectively

$$(A) \quad \sum_{i=1,2} |U_{ei}|^2 \leq a_e^0, \quad \sum_{i=3,4} |U_{\mu i}|^2 \leq a_\mu^0, \quad (21)$$

$$(B) \quad \sum_{i=3,4} |U_{ei}|^2 \leq a_e^0, \quad \sum_{i=1,2} |U_{\mu i}|^2 \leq a_\mu^0. \quad (22)$$

For the amplitude of $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ oscillations in both schemes we have the upper bound

$$A_{\mu;e} \leq 4 \text{Min}(a_e^0, a_\mu^0), \quad (23)$$

which is compatible with the LSND result.

If scheme A is realized in nature, the experiments on the measurement of the neutrino mass with the ^3H method and the experiments on the search for neutrinoless double-beta decay have a good chance to see the effects of the SBL mass-squared difference Δm^2 . In fact, taking into account the inequalities (21), for the mass $m(^3\text{H})$ and for the effective Majorana mass $\langle m \rangle = \sum_{i=1}^n U_{ei}^2 m_i$ in scheme A we have

$$m(^3\text{H}) \simeq \sqrt{\Delta m^2}, \quad (24)$$

$$|\langle m \rangle| \simeq \left| \sum_{i=3,4} U_{ei}^2 \right| \sqrt{\Delta m^2} \leq \sqrt{\Delta m^2}. \quad (25)$$

In the framework of the schemes A and B, rather strong constraints for the probabilities of long-baseline (LBL) $\bar{\nu}_e \rightarrow \bar{\nu}_e$ and $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ transitions can be inferred from the bounds on the elements of the mixing matrix obtained from SBL experiments [4]. Let us consider scheme A. For the probabilities of $\nu_\alpha \rightarrow \nu_\beta$ transitions in LBL experiments we have

$$P_{\nu_\alpha \rightarrow \nu_\beta}^{(\text{LBL}, \text{A})} = \left| \sum_{k=1,2} U_{\beta k} U_{\alpha k}^* e^{-i \frac{\Delta m_{k1}^2 L}{2p}} \right|^2 + \left| \sum_{k=3,4} U_{\beta k} U_{\alpha k}^* \right|^2. \quad (26)$$

The corresponding equation for antineutrinos implies that the probability of LBL $\bar{\nu}_e \rightarrow \bar{\nu}_e$ transitions is bounded by

$$P_{\bar{\nu}_e \rightarrow \bar{\nu}_e}^{(\text{LBL})} \geq \left(1 - \sum_{i=1,2} |U_{ei}|^2 \right)^2. \quad (27)$$

From the results of SBL reactor experiments it follows that $\sum_{i=1,2} |U_{ei}|^2$ is small (see Eq.(21)) and thus (in the scheme under consideration) the LBL probability $P_{\bar{\nu}_e \rightarrow \bar{\nu}_e}^{(\text{LBL})}$ must be close to one. In




Figure 2

fact, from Eqs.(21) and (27), for the probability of the LBL transitions of $\bar{\nu}_e$'s into all possible states we have the upper bound

$$1 - P_{\bar{\nu}_e \rightarrow \bar{\nu}_e}^{(LBL)} \leq a_e^0 (2 - a_e^0) . \quad (28)$$

The same inequality is valid in scheme B.

The upper bound (28) is presented in Fig.2 for the values of the SBL parameter Δm^2 in the range (8) (the solid line). The dash-dotted and dash-dot-dotted lines correspond to the expected sensitivity of the CHOOZ and Palo Verde LBL reactor experiments [12]. As it is seen from Fig.2, the sensitivity of the CHOOZ experiment could allow to see effect of neutrino oscillations only if $\Delta m^2 \gtrsim 3 \text{ eV}^2$. The shadowed region in Fig.2 corresponds to the range of Δm^2 allowed by the results of the LSND experiment, taking into account the results of all the other SBL experiments.

In conclusion, we have considered here the following question: what information about the neutrino mass spectrum and the elements of the neutrino mixing matrix can be inferred from the results of neutrino oscillation experiments? We came to the conclusion that from the existing data it follows that the general characteristics of neutrino and quark mixing are quite different. Only future neutrino oscillation experiments will allow to obtain information about the genuine neutrino mass spectrum and the neutrino mixing matrix.

REFERENCES

1. S.M. Bilenky *et al.*, Phys. Lett. B **356**, 273 (1995); Phys. Rev. D **54**, 1881 (1996).
2. S.M. Bilenky *et al.*, Phys. Rev. D **54**, 4432 (1996).
3. S.M. Bilenky *et al.*, hep-ph/9607372.
4. S.M. Bilenky *et al.*, hep-ph/9705300.
5. B. Pontecorvo, J. Exptl. Theoret. Phys. **33**, 549 (1957) [Sov. Phys. JETP **6**, 429 (1958)]; *ibid.* **34**, 247 (1958) [*ibid.* **7**, 172 (1958)].
6. B.T. Cleveland *et al.*, Nucl. Phys. B (Proc. Suppl.) **38**, 47 (1995); K.S. Hirata *et al.*, Phys. Rev. D **44**, 2241 (1991); GALLEX Coll., Phys. Lett. B **388**, 384 (1996); J.N. Abdurashitov *et al.*, Phys. Rev. Lett. **77**, 4708 (1996).
7. Y. Fukuda *et al.*, Phys. Lett. B **335**, 237 (1994); R. Becker-Szendy *et al.*, Nucl. Phys. B (Proc. Suppl.) **38**, 331 (1995); W.W.M. Allison *et al.*, Phys. Lett. B **391**, 491 (1997).
8. C. Athanassopoulos *et al.*, Phys. Rev. Lett. **77**, 3082 (1996).
9. B. Achkar *et al.*, Nucl. Phys. B **434**, 503 (1995).
10. F. Dydak *et al.*, Phys. Lett. B **134**, 281 (1984); I.E. Stockdale *et al.*, Phys. Rev. Lett. **52**, 1384 (1984).
11. X. Shi and D.N. Schramm, Phys. Lett. B **283**, 305 (1992).
12. R.I. Steinberg, Proc. of the 5th Int. Workshop on Neutrino Telescopes, Venezia, March 1993; F. Boehm *et al.*, The Palo Verde experiment, 1996.